

Instructions for Avian Influenza (Subtype H9) Vaccine, Inactivated (Strain NJ01)

NAME OF VETERINARY MEDICINE

English name: Avian Influenza (Subtype H9) Vaccine, Inactivated (Strain NJ01)

Chinese pinyin: Qinliugan (H9 Yaxing) Miehuoyimiao (NJ01 Zhu)

MAIN INGREDIENTS AND CONTENT

This product contains inactivated H9 subtype avian influenza NJ01 strain virus, the virus content before inactivation is not less than $10^{8.0} \text{ELD}_{50}/0.1\text{ml}$.

CHARACTER

Milky white emulsion

FUNCTIONS AND USES

It is used to prevent avian influenza in ducks and chickens caused by H9 subtype avian influenza virus.

Ducks : One immunization: 8 ~ 12 days of age, the immunization period was 3 weeks. The age of immunization was 3 ~ 5 weeks, and the immunization period was 3 months. Twice immunization: first immunization at 8-12 days of age, second immunization at 5 weeks of age, and the immunization period was 5 months.

Chicken: primary immunization: 10 ~ 12 days old, the immunization period was 6 weeks. The age of immunization was 3 ~ 5 weeks, and the immunization period was 4 months. Twice immunization: first immunization at 10-12 days of age, second immunization at 7 weeks of age, and the immunization period was 6 months.

USAGE AND DOSAGE

Muscle or neck back subcutaneous injection.

Ducks : under 5 weeks of age, each 0.3ml ; above 5 weeks of age, each 0.5ml; 1 ~ 2 weeks prior to the start of egg production, each 0.8ml : after every 5 months to strengthen the immunization, each 0.8ml.

Chicken : less than 5 weeks old, each 0.3ml ; above 5 weeks of age, each 0.5ml ; after that, the immunization was strengthened every 6 months, each. 0.5 ml.

ADVERSE REACTIONS

Generally no adverse reactions.

MATTERS NEEDING ATTENTION

- (1) Only for vaccination of healthy chickens and ducks.
- (2) Prevent freezing and high temperature.
- (3) The vaccine should reach room temperature and shake well before use.
- (4) Prior to use, the vaccine should be carefully examined. If abnormalities such as demulsification or deterioration are found, it must not be used.
- (5) During vaccination, local disinfection should be carried out.
- (6) After the vaccine bottle is opened, it should be used up on the same day.
- (7) It is forbidden to be used for meat ducks and broiler chickens within 21 days before slaughter.

SPECIFICATIONS

(1) 100 dose / bottle (2) 250 dose / bottle (3) 500 dose / bottle

PACKAGE

(1) 100 bottles/box (2) 40 bottles/box (3) 20 bottles/box

STORAGE AND STORAGE PERIOD

Stored below 2~8°C, the validity period is 15 months.

APPROVAL NUMBER

SYSZ240052217

PRODUCTION ENTERPRISE

Guizhou Firstv Biological Technology Co.Ltd

Address: No. 268, kechuang south road, Baiyun District, Guiyang City, Guizhou Province, China.

Postcode: 550014

The phone: 0851-84603559 Fax: 0851-84603559



USE ONLY UNDER VETERINARY GUIDANCE