

***Newcastle Disease, Avian Influenza (Subtype H9) and Fowl Adenoviruses
Disease (Group I, Serotype 4) Vaccine,
Inactivated (Strain La Sota+ Strain HS+Strain DN)***

NAME OF VETERINARY MEDICINE

English name: Newcastle Disease, Avian Influenza (Subtype H9) and Fowl Adenoviruses Disease
(Group I, Serotype 4) Vaccine, Inactivated (Strain La Sota+ Strain HS+Strain DN)
Chinese pinyin: JiXinchengyi, Qinliugan (H9Yaxing), Qinxianbingdubing (I Qun, 4Xing) Sanlian
Miehuoyimiao (La Sota Zhu+HS Zhu+DN Zhu)

MAIN INGREDIENTS AND CONTENT

The vaccine contains inactivated chicken Newcastle disease virus La Sota strain (the virus content before inactivation $\geq 10^{8.0}$ EID₅₀/0.1ml), and the Avian influenza virus (H9 subtype) HS strain (the virus content before inactivation $\geq 10^{8.0}$ EID₅₀/0.1 ml), Avian adenovirus (group I, serotype 4) DN strain (the virus content before inactivation $\geq 10^{7.0}$ TCID₅₀/0.1ml).

CHARACTER

Milky white homogeneous emulsion.

FUNCTIONS AND USES

It is used to prevent Newcastle Disease, Avian Influenza (H9 subtype) and disease caused by avian adenovirus (group I, serotype 4). The immunization period is 4 months for 7 ~ 21 days old chickens and 6 months for 21 days of age and above chickens.

USAGE AND DOSAGE

Subcutaneous or intramuscular injection in the neck, 0.3ml for chickens within 7~21 days of age, 0.5ml for chickens 21 days of age and above.

ADVERSE REACTIONS

Generally, there are no visible adverse reactions.

MATTERS NEEDING ATTENTION

- (1) This vaccine should only be administered to healthy chickens.
- (2) Before use, the vaccine should be carefully inspected, and should not be used if it is found to be discolored, demulsification, broken or leaking, or mixed with foreign matter.
- (3) Before use, the vaccine should be restored to room temperature and shaken well.
- (4) Vaccination instruments should be sterile and the injection site should be sterilized.
- (5) After the vaccine is opened, it should be used up on the same day.
- (6) Unused vaccines, used vaccine vials and syringes should be disposed of harmlessly.
- (7) Vaccines should not be frozen during transportation and storage, and the use of frozen vaccines is strictly prohibited.
- (8) Prohibit the use of vaccines within 21 days before slaughter.

STORAGE AND STORAGE PERIOD

Store at 2~8℃, and the validity period is 18 months.

APPROVAL NUMBER

SYSZ 240052392

PRODUCTION ENTERPRISE

Guizhou Firstv Biological Technology Co.Ltd

Address: No. 268, kechuang south road, Baiyun District, Guiyang City, Guizhou Province, China.

Postcode: 550014

The phone: 0851-84603559 Fax: 0851-84603559

USE ONLY UNDER VETERINARY GUIDANCE

Firstv[®]